



Task Title: Medicine Label

OALCF Cover Sheet – Learner Copy

Learner Name: _____

Date Started: _____

Date Completed: _____

Successful Completion: Yes No

Goal Path: Employment Apprenticeship

Secondary School Post Secondary Independence

Task Description: Read a medicine label and interpret directions and correct dosage.

Main Competency/Task Group/Level Indicator:

- Find and Use Information/Interpret documents/A2.2
- Understand and Use Numbers/Manage time/C2.1
- Understand and Use Numbers/Use measures/C3.1

Materials Required:

- Pen/pencil and paper and/or digital device
- Calculator or digital device with calculator function (optional)

Learner Information

Many people need to accurately read and dispense medication for their work and in their daily lives.

Scan the "Adult Robitussin" cough medicine label.

**Packaged with Tamper-Evident bottle cap.
Do Not Use if breakable ring is separated or missing.**

Active ingredient (in each 10 ml):
Guaifenesin, USP 200 mg

Use: helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive.

Warnings
Ask a doctor before use if you have cough that occurs with too much phlegm (mucus); cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema.
Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Other information: each 10 ml contains: sodium 4 mg. Store at 20-25°C (68-77°F). Alcohol-free.

ADULT

Robitussin®

Mucus + Chest Congestion

GUAIFENESIN (Expectorant)

Alcohol-Free

4 FL OZ (118 ml)

Inactive ingredients: anhydrous citric acid, artificial flavor, caramel, FD&C red no. 40, glycerin, high fructose corn syrup, liquid glucose, menthol, propylene glycol, purified water, saccharin sodium, sodium benzoate

Questions or comments?
Call weekdays from 9 AM to 5 PM EST 1-800-762-4675

Directions: do not take more than 6 doses in any 24-hour period. Measure only with dosing cup provided. Keep dosing cup with product. ml = milliliter. This adult product is not intended for use in children under 12 years of age.

age	dose
adults and children 12 years and over	10 – 20 ml every 4 hours
children under 12 years	do not use

Distributed by:
Pfizer, Madison, NJ 07940 USA
© 2012 Pfizer Inc.

For most recent product information, visit www.robitussin.com

LOT:
EXP:

PAA035209

Work Sheet

Task 1: What is this medication used for?

Answer:

Task 2: When should someone see a doctor or seek advice from a healthcare professional?

Answer:

Task 3: At what temperature should this medicine be stored?

Answer:

Task 4: What is the maximum dosage for an adult in a 24-hour period?

Answer:

Task 5: What is the maximum dosage for a child under 12 in a 24-hour period?

Answer:

Task Title: MedicineLabel_EI_A2.2_C2.1_C3.1

Task 6: If you give someone a dose of this medication at 10:30am, what is the earliest time you can give another dose? Assume the person has not yet reached the maximum dosage within a 24-hour period.

Answer:
